

風治散

THIS MEDICINE FOR GENERAL USE
FUUJISAN (Kakkonto Extract Granules)
(Category 2 OTC medicines)

INDICATIONS :

Following symptoms in person with moderate or more physical strength; Initial stage of common cold (without spontaneous sweating), coryza, rhinitis, headache, shoulder stiffness, muscle pain, and/or pain in hands and shoulder.

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of strength or constitutional weakness).
 - (4) Persons who have gastrointestinal weakness.
 - (5) Persons with excessive sweating.
 - (6) Elderly persons.
 - (7) Persons who have experienced rash • redness, itching. etc. due to medicines or something.
 - (8) Persons with the following symptoms: edema, difficulty in urination.
 - (9) Persons diagnosed as having the following: hypertension, heart disease, kidney disease, thyroid disease.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash • redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately

Symptom's name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually worsen.
Hepatic function failure	Following symptoms appear: fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month (5 to 6 times if you take this medicine for in the case of the initial stage of the common cold, coryza, and headache). Take this leaflet with you.
4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following dosages with warm water or water, before or after meals.

Age	One dosage	Daily dosage
Adult (15 years or over)	1 Packet (2.5g)	3 times
7 years or over ~ under 15 years	2/3 Packet	
4 years or over ~ under 7 years	1/2 Packet	
2 years or over ~ under 4 years	1/3 Packet	
Less than 2 years	Do not use	

CAUTION ASSOCIATED WITH DOSAGE

This medication should be given to children only under adult supervision.

INGREDIENTS

3 Packets

Composition	Dosage	
Kakkon (Pueraria Root)	4.0g	2.8g of a water extract of the mixed crude drugs.
Maou (Ephedra Herb)	3.0g	
Shokyo (Ginger)	1.0g	
Taiso (Jujube)	3.0g	
Keihi (Cinnamon Bark)	2.0g	
Shakuyaku (Peony Root)	2.0g	
Kanzo (Glycyrrhiza)	2.0g	

Inactive ingredients: starch

CAUTION

Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKAN-YAKU MEDICAL INSTITUTE CO.LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)