

ササイサン

THIS MEDICINE FOR GENERAL USE

SASAI SAN (Granule)

(Category 2 OTC medicines)

INDICATIONS :

Loss of appetite, stomach and abdominal bloating, dyspepsia, gastrointestinal weakness, overeating, overdrinking, heartburn, heavy stomach, the pressure on the chest, nausea (including retching, upset stomach, sick feeling by hangover and/or sick from drinking), vomiting

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Elderly persons.
 - (3) Persons with the following symptom; edema.
 - (4) Persons diagnosed as having the following; hypertension, kidney disease, heart disease.
2. In very rare cases the following serious symptoms may occur after taking medicine. In this case, consult a physician immediately.

Symptom's name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, stretched feelings, and stiffness of hands and feet, feeling of weakness and muscular pain with appear and gradually worsen.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month. Take this leaflet with you.

4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following dosages with warm or water, between meals.

(“Between meals” means a couple of hours after meals.)

Age	One dosage	Daily dosage
Adult (15 years or over)	1 Packet (1.5g)	3 times
11 years or over ~ under 15 years	2/3 Packet	
8 years or over ~ under 11 years	1/2 Packet	
5 years or over ~ under 8 years	1/3 Packet	
3 years or over ~ under 5 years	1/4 Packet	
Less than 3 years	Do not use	

CAUTION ASSOCIATED WITH DOSAGE

The medication should be given to children only under adult supervision.

INGREDIENTS

3 Packet

Composition	Dosage
Kanzo (Glycyrrhiza) extract (Equivalent to 3.6g of raw Glycyrrhiza)	0.9g
Oubaku (Phellodendron Bark) extract (Equivalent to 2.5g of raw Phellodendron Bark)	0.4g
Hakka (Mentha Herb)	0.3g

Inactive ingredients: lactose hydrate

CAUTION

Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.

- (2) Keep all medication out of the reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKAN-YAKU MEDICAL INSTITUTE CO. LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)