

紫華栄

THIS MEDICINE FOR GENERAL USE

SHIKARON (Granule)

(Category 3 OTC medicines)

INDICATIONS

Revitalization in persons with following conditions; Weak constitution, physical fatigue, weakness during and after illness, gastrointestinal weakness, loss of appetite, pale complexion, over-sensitive to cold, and revitalization in growth season.

CAUTION FOR USE

• Who should seek consultation

1. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash・redness, itching

2. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following dosages with warm water or water, between meals.

(“Between meals” means a couple of hours after meals.)

Age	One dosage	Daily dosage
Adult (15 years or over)	1 Packet (1.5g)	3 times
11 years~under 15 years	1/2 Packet	
8 years~under 11 years	1/3 Packet	

CAUTION ASSOCIATED WITH DOSAGE

- (1) Observe the dosage and usage.
- (2) The medication should be given to children only under adult supervision.

INGREDIENT

3 Packet

Composition	Raw material	Dosage
Shikon (Lithospermum Root) extract	7,500mg	170mg
Touki (Japanese Angelica Root) extract	342mg	120mg
Zyuyaku (Houttuynia Herb) extract	1,500mg	85mg
Ninjin (Ginseng) extract	3,000mg	600mg
Ougi (Astragalus Root) extract	250mg	50mg
Senkyu (Cnidium Rhizome) extract	225mg	50mg
Yokuinin powder (Powdered Coix Seed)		1,800mg
Kanzo powder (Powdered Glycyrrhiza)		45mg

Inactive ingredients: lactose hydrate

CAUTION

Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKAN-YAKU MEDICAL INSTITUTE CO. LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)