

松寿仙

THIS MEDICINE FOR GENERAL USE

SYOJUSEN

INDICATIONS :

Revitalization in the following cases; weak constitution, physical exsaustion, during and/or after sickness, gastrointestinal weakness, loss of appetite.

REACTIONS

1. Please consult doctors, pharmacists or registered distributors with the following cases, before taking this medicine.
 - (1) Having a medication from medical doctors.
 - (2) Individual and family have allergies.
 - (3) Having a history of developing allergic symptoms after taking medicine, etc.
2. If you experience any of the following symptoms which are possibly side effects, adminisitation should be discontinued immediately and consult doctors, pharmacists or registered distributors with this package incert.

Affected parts	Symptoms
Skin	Rash, redness, itch

3. Consult doctors, pharmacists, or registered distributors if you want to use this product for a long period of time.

DOSES AND ADMINISTRATION

Take the adequate dosage after dilution with warm or cold water, before or between meals.

(“Between meals” means a couple of hours after meals.)

Age	One dosage	one day
Adult (over 15 years old)	2ml	3 times
15~8 years old	1ml	

CAUTION ASSOCIATED WITH DOSES AND ADMINISTRATION

- (1) Please observe the direction and doses.
- (2) In case of infants, use it under guardian's direction.

COMPOSITION

Composition (in 6ml)	Dosage (in 6ml)
Chlorophyllous liquid of Sasa (Equivalent to 19g of raw Sasa)	4.8ml
Akamatsu-ba extract (Japanese red pine extract) (Equivalent to 232mg of raw Japanese red pine)	18.0mg
Ninjin extract (Ginseng extract) (Equivalent to 58mg of raw Ginseng)	10.8mg

Inactive ingredients : paraben

CAUTION

Since the medicine contains crude drugs, the liquid color may vary between individual products. However this has no influence on the efficacy of the medicine.

DIRECTIONS FOR HANDLING AND KEEPING

- (1) To keep away from reaching children.
- (2) To keep it in a dark and dry cool place.
- (3) Not to touch the spout directly when opening and taking to avoid a quality variation by a contamination from hands.
- (4) Be careful not to cut fingers when opening and separating off a plastic loop.
- (5) To keep it in refrigerator with putting cap back on the bottle.
- (6) Content fluid may spills in case of carrying the bottle with the part of cap and/or keeping the bottle sideways, under a condition of failing to close the cap tightly.
- (7) Content fluid may transform depending on how to use of it. To follow precautions strictly after breaking the seal.
- (8) Do not replace into other containers to avoid a misuse and secure the quality.
- (9) Since the medicine contains crude drugs, amino acid and the like may attach to the side or the bottom of the bottle. However this has no influence on the efficacy of the medicine by taking after shaking the bottle.

MANUFACTURED AND DISTRIBUTED BY :
WAKAN-YAKU MEDICAL INSTITUTE CO.,LTD.
1-29-8, Shinjuku, Shinjuku-ku, Tokyu, Japan