

新ササカル

THIS MEDICINE FOR GENERAL USE

SHIN SASA-CAL (Tablet)

(Category 3 OTC medicines)

INDICATIONS :

a) Growth promotion of bones and/or teeth in the following cases: weak constitution, scrofulosis*

*scrofulosis means frail or helpless constitution of persons having a tendency to get anemia etc.

b) Prevention of weakness of bones and/or teeth during pregnancy or lactation

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

Persons undergoing medical treatment from a physician.

2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash • redness, itching

3. The following symptom may appear after taking this medicine. If the symptom continues or worsen, stop taking and consult a physician, pharmacist or registered distributor. Take this leaflet with you.: constipation

4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following dosages with warm water or water, between meals.

(“Between meals” means a couple of hours after meals.)

Age	One dosage	Daily dosage
15 years or over	5 Tablets	3 times
15~7 years	3 Tablets	
7~5 years	2 Tablets	
Less than 5 years	Do not use	

CAUTION ASSOCIATED WITH DOSAGE

- (1) Observe the dosage and usage.
- (2) This medication should be given to children only under adult supervision.

INGREDIENTS

15 Tablets

Composition	Dosage	Behavior
Borei powder (Powdered Oyster Shell)	1,500mg (Containing 470mg of Calcium)	Calcium supplements
Calcium Gluconate Hydrate	800mg (Containing 70mg of Calcium)	Calcium supplements
Calcium Lactate Hydrate	500mg (Containing 60mg of Calcium)	Calcium supplements
Dried Yeast	300mg	Supplements of vitamin Bs, amino acids, etc., and increasing calcium absorption

Inactive ingredients: starch, lactose hydrate, sodium copper chlorophyllin, magnesium stearate, flavor

CAUTION

Since the medicine contains crude drugs, the tablet color may vary for individual products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store hermetically in a cool dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) Do not take a desiccant in the package.

MANUFACTURED AND DISTRIBUTED BY:

WAKAN-YAKU MEDICAL INSTITUTE CO. LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)