

和漢薬ササラック

THIS MEDICINE FOR GENERAL USE

WAKAN-YAKU SASA-LUCK (Granule)

(Category designated ② OTC medicines)

INDICATIONS :

Constipation, following symptoms associated with constipation: heavy head, hot flashes, rough skin, pimple, loss of appetite, abdominal bloating, abnormal intestinal fermentation and hemorrhoid.

CAUTION FOR USE

• TO BE AVOIDED

FAILURE TO OBSERVE THE FOLLOWING MAY RESULT IN THE SYMPTOMS WORSENING OR THE OCCURRENCE OF ADVERSE REACTIONS/ACCIDENTS.

1. THIS MEDICATION SHOULD NOT BE USED IN COMBINATION WITH THE FOLLOWING MEDICATIONS.

OTHER CATHARTIC AGENTS

2. WOMEN WHO ARE NURSING SHOULD NOT TAKE THIS MEDICINE. IF THOSE WOMEN TAKE IT, AVOID BREAST FEEDING.

3. DO NOT TAKE IN LARGE QUANTITIES.

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

(1) Persons undergoing medical treatment from a physician.

(2) Pregnant women or women suspected of being pregnant.

(3) Persons who are physically weak(declining of strength or constitutional weakness).

(4) Persons with weak stomach and intestines who have a tendency to diarrhea.

(5) Persons who have experienced allergic symptom due to medicines or something.

(6) Persons with the following symptoms: severe abdominal pain, nausea, vomiting.

2. The following symptoms may be adverse reactions to this medication. If any of

these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash • redness, itching
Gastrointestinal system	Severe abdominal pain, nausea • vomiting.

3. The following symptom may appear after taking this medicine. If the symptom continues or worsen, stop taking and consult a physician, pharmacist or registered distributor. Take this leaflet with you:

Diarrhea

4. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 week. Take this leaflet with you.

DOSAGE

Take the medicine on an empty stomach 1 or 3 times a day, up to 3 times a day.

Wait more than four hours between doses.

Start from minimal dose and then increase or decrease until appropriate dose for the state or condition of the bowel movements.

Age	One dosage	Daily dosage
15 years or over	3/4 ~ 1 Packet	1 ~ 3 times
11 years or over ~ under 15 years	1/2 ~ 2/3 Packet	
7 years or over ~ under 11 years	2/5 ~ 1/2 Packet	
3 years or over ~ under 7 years	1/4 ~ 1/3 Packet	
Less than 3 years	Do not use	

CAUTION ASSOCIATED WITH DOSAGE

- (1) Observe the dosage and usage.
- (2) This medication should be given to children only under adult supervision.
- (3) When children who are 3 years or over take this medication, be careful it does not get stuck in the throat.

INGREDIENTS

Composition	Daily dosage 4.5g (3 Packet)
Daiou (Powdered Rhubarb)	1.0g
Senna (Powdered Senna Leaf)	1.0g
Kanzo (Powdered Glycyrrhiza)	0.6g
Shakuyaku (Powdered Peony Root)	0.5g
Hakka (Powdered Mentha Herb)	0.5g

Inactive ingredients : corn starch

CAUTION

Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store hermetically in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY :

WAKAN-YAKU MEDICAL INSTITUTE CO.,LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)