和漢薬晴快散

THIS MEDICINE FOR GENERAL USE WAKAN-YAKU SEIKAISAN (Granule) (Category 3 OTC medicines)

INDICATIONS :

Forgetfulness after middle age

CAUTION FOR USE

 \cdot To be avoided

Failure to observe the following may result in the symptoms worsening or the occurrence of adverse reactions/accidents.

The following persons do not use.

- (1) Persons diagnosed as having the following: dementia
- (2) Persons suspected of being dementia
- $\boldsymbol{\cdot}$ Who should seek consultation
- 1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who have experienced allegic symptom, etc. due to medicines or something
- 2. In case of the following, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.
- (1) the following symptoms occur, after taking this medicine.

Related areas	Symptoms	
Skin	Rash \cdot redness, itching	
Gastrointestinal system	Nausea \cdot vomiting, loss of appetite, gastric	
	distress	
Neuropsychiatric system	Dizziness	

- 3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month. Take this leaflet with you.
- 4. The following symptom may occur after taking medicine: Diarrhea. If the continuance or enhancement of such a symptom is observed, stop taking this medicine and consult a physician, pharmacist or registered distributor. Take this leaflet with you.
- 5. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

OTHER CAUTIONS

Taking this medicine may influence the laboratory value (1,5-AG in the blood) regarding diabetes.

DOSAGE

Take the following dosages with warm water or water, before or between meals.

Age	One dosage	Daily dosage
Adult (15 years or over)	1 Packet (1.2g)	3 times
Less than 15 years	Do not use	

("Between meals" means a couple of hours after meals.)

INGREDIENTS

3 Packets

Composition	Dosage
Onji (Polygara Root) extract	1.3g
(Equivalent to 3g of raw Polygara Root)	

Inactive ingredients: dextrin, magnesium stearate, light anhydrous silicic acid, crystalline cellulose, lactose hydrate

PRECAUTIONS FOR STORAGE AND HANDLING

(1) Store in a cool, dry place free from direct sunlight.

(2) Keep all medication out of the reach of children.

- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.

MANUFACTURED AND DISTRIBUTED BY: WAKAN-YAKU MEDICAL INSTITUTE CO. LTD. 1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)