松寿仙

# THIS MEDICINE FOR GENERAL USE SYOJUSEN (liquid) (Category 3 OTC medicines)

## **INDICATIONS** :

Revitalization in persons with following conditions; weak constitution, physical fatigue, during and after illness, gastrointestinal weakness, loss of appetite.

#### CAUTION FOR USE

- $\boldsymbol{\cdot}$  Who should seek consultation
- 1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - (1) Persons undergoing medical treatment from a physician.
  - (2) Persons who are allergic in person or family.
  - (3) Persons who have experienced allergic symptoms due to medicines or something.
- 2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms	
Skin	Rash•redness, itching	

3. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

## DOSAGE

Take the following dosages measured with the attached measuring cup, after dilution with warm water or water, before or between meals.

("Between meals" means a couple of hours after meals.)

Age	One dosage	Daily dosage	
Adult (15 years or over)	2ml	2ml 1ml 3 times	
8 years~under 15 years	1ml		

# CAUTION ASSOCIATED WITH DOSAGE

(1) Observe the dosage and usage.

(2) The medication should be given to children only under adult supervision.

# INGREDIENT

Composition (in 6ml)	Dosage (in 6ml)
Chlorophyllous liquid of Sasa	4.8ml
(Equivalent to 19g of raw Sasa)	
Akamatsu-ba extract (Japanese red pine extract)	18.0mg
(Equivalent to 232mg of raw Japanese red pine)	
Ninjin extract (Ginseng extract)	10.8mg
(Equivalent to 58mg of raw Ginseng)	

Inactive ingredients: paraben

## CAUTION

Since the medicine contains crude drugs, the liquid color may vary for individual products, however this has no influence on the efficacy of the medicine.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep all medication out of the reach of children.
- (2) Store in a cool place free from direct sunlight.
- (3) Not to touch the spout directly when opening and taking to prevent quality deterioration by a contamination from hands.
- (4) Be careful not to cut fingers when opening and separating off a plastic loop.
- (5) Store in refrigerator with putting cap back on the bottle after opening, and use up as early as possible.
- (6) After opening, content fluid may spill in case of carrying the bottle with the part of cap and/or keeping the bottle sideways under a condition of failing to close the cap tightly.

- (7) Do not transfer to another container to prevent misuse and quality deterioration.
- (8) Since the medicine is herbal extract formulation, ingredient such as amino acid may attach to the side or the bottom of the bottle, however this has no influence on the efficacy of the medicine by taking after shaking the bottle.

# MANUFACTURED AND DISTRIBUTED BY: WAKAN-YAKU MEDICAL INSTITUTE CO. LTD. 1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: December 2023 (Second edition)